arXiv:1511.04983v1 [math.CO] 16 Nov 2015

Distance integral complete multipartite graphs with $s = 5, 6$ *†

Ruosong Yang, Ligong Wang‡
Department of Applied Mathematics, School of Science, Northwestern Polytechnical University,
Xi'an, Shaanxi 710072, People's Republic of China.
E-mail: lnxyangruosong@163.com, lgwangmath@163.com

Abstract

Let $D(G) = (d_{ij})_{n\times n}$ denote the distance matrix of a connected graph G with order n, where d_{ij} is equal to the distance between vertices v_i and v_j in G. A graph is called distance integral if all eigenvalues of its distance matrix are integers. In 2014, Yang and Wang gave a sufficient and necessary condition for complete r-partite graphs $K_{p_1,p_2,\ldots,p_r} = K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$ to be distance integral and obtained such distance integral graphs with $s = 1, 2, 3, 4$. However distance integral complete multipartite graphs $K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$ with $s > 4$ have not been found. In this paper, we find and construct some infinite classes of these distance integral graphs $K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$ with $s = 5, 6$. The problem of the existence of such distance integral graphs $K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$ with arbitrarily large number s remains open.

Key Words: Complete multipartite graph, Distance matrix, Distance integral, Graph spectrum.

2000 MR Subject Classification: 05C50, 05C12.

1 Introduction

Let G be a simple connected undirected graph with vertex set denoted by $V(G) = \{v_1, v_2, \ldots, v_n\}$. $A(G) = (a_{ij})$ is an $n \times n$ matrix, where $a_{ij} = 1$ if v_i and v_j are adjacent and $a_{ij} = 0$ otherwise. The distance between the vertices v_i and v_j is the length of a shortest path between them, and is denoted by d_{ij}. The distance matrix of G, denoted by $D(G)$, is the $n \times n$ matrix whose (i, j)-entry is equal to d_{ij} for $i, j = 1, 2, \ldots, n$ (see [3]). Note that $d_{ii} = 0$, $i = 1, 2, \ldots, n$. The distance characteristic polynomial (or D-polynomial) of G is $D_G(x) = |xI_n - D(G)|$, where I_n is the $n \times n$ identity matrix. The eigenvalues of $D(G)$ are said to be the distance eigenvalues or D-eigenvalues of G. Since $D(G)$ is a real symmetric matrix, the D-eigenvalues are real and can be denoted as $\mu_1 \geq \mu_2 \geq \ldots \geq \mu_n$. The distance spectral radius of G is the largest D-eigenvalue μ_1 and denoted by $\mu(G)$. Assume that $\mu_1 > \mu_2 > \ldots > \mu_t$ are t distinct D-eigenvalues of G with the corresponding multiplicities $k_1, k_2, \ldots, k_t$. We denote by $Spec(G) = \begin{pmatrix} \mu_t & \mu_{t-1} & \cdots & \mu_2 & \mu_1 \\ k_t & k_{t-1} & \ldots & k_2 & k_1 \end{pmatrix}$ the Distance spectrum or the D-spectrum of G. G is called integral if all eigenvalues of $A(G)$ are integers. Similarly to integral graphs, a graph is called distance integral if all its D-eigenvalues are integers. Many results about distance spectral radius and the D-eigenvalues of graphs can be found in [1, 2, 4, 6, 7, 9, 11, 12, 15, 17, 18]. Some more results about spectral graph theory can be found in [8, 13, 19].

*Supported by the National Natural Science Foundation of China (No.11171273) and graduate starting seed fund of Northwestern Polytechnical University (No.Z2014173).

†Ruosong Yang (1990-), male, native of LuoYang, Henan, an postgraduate student of Northwestern Polytechnical University, engages in graph theory and its application.

‡Corresponding author

A *complete r-partite* ($r \geq 2$) *graph* $K_{p_1,p_2,\cdots,p_r}$ is a graph with a set $V = V_1 \cup V_2 \cup \cdots \cup V_r$ of $p_1 + p_2 + \cdots + p_r (= n)$ vertices, where V_i's are nonempty disjoint set, $|V_i| = p_i$, such that two vertices in V are adjacent if and only if they belong to different V_i's. Assume that the number of distinct integers of $p_1, p_2, \cdots, p_r$ is s. Without loss of generality, assume that the first s ones are the distinct integers such that $p_1 < p_2 < \ldots < p_s$. Suppose that a_i is the multiplicity of p_i for each $i = 1, 2, \ldots, s$. The complete r-partite graph $K_{p_1,p_2,\cdots,p_r} = K_{p_1,\ldots,p_1,\ldots,p_s,\ldots,p_s}$ on n vertices is also denoted by $K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$, where $r = \sum_{i=1}^{s} a_i$ and $n = \sum_{i=1}^{s} a_i p_i$. In 2014, Yang and Wang gave a sufficient and necessary condition for $K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$ to be distance integral and obtained such distance integral graphs $K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$ with $s = 1, 2, 3, 4$ (see[16]). In this paper, we find and construct some infinite classes of these distance integral graphs $K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$ with $s = 5, 6$ by using a computer search. For $s = 5, 6$, we give a positive answer to a question of Yang et al. [16]. The problem of the existence of distance integral complete multipartite graphs $K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$ with arbitrarily large number s remains open.

2 Preliminaries

In this section, we shall give some known results on distance integral complete r-partite graphs, which are used in our theorem proof later.

Theorem 1. *(See Theorem 4.1 of [7] and [10]) Let G be a complete r-partite graph $K_{p_1,p_2,\ldots,p_r}$ on n vertices. Then the D-polynomial of G is*

$$D_G(x) = \prod_{i=1}^{r}(x+2)^{(p_i-1)} \prod_{i=1}^{r}(x - p_i + 2)(1 - \sum_{i=1}^{r} \frac{p_i}{x - p_i + 2}). \tag{1}$$

Theorem 2. *(See [16]) If the complete r-partite graph $K_{p_1,p_2,\ldots,p_r} = K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$ on n vertices is distance integral then there exist integers $\mu_i (i = 1, 2, \ldots, s)$ such that $-2 < p_1 - 2 < \mu_1 < p_2 - 2 < \mu_2 < \cdots < p_{s-1} - 2 < \mu_{s-1} < p_s - 2 < \mu_s < +\infty$ and the numbers $a_1, a_2, \ldots a_s$ defined by*

$$a_k = \frac{\prod_{i=1}^{s}(\mu_i - p_k + 2)}{p_k \prod_{i=1, i\neq k}^{s}(p_i - p_k)}, k = 1, 2, \ldots, s, \tag{2}$$

are positive integers.

Conversely, suppose that there exist integers μ_i $(i = 1, 2, \ldots, s)$ such that $-2 < p_1 - 2 < \mu_1 < p_2 - 2 < \mu_2 < \cdots < p_{s-1} - 2 < \mu_{s-1} < p_s - 2 < \mu_s < +\infty$ and that the numbers $a_k = \frac{\prod_{i=1}^{s}(\mu_i - p_k + 2)}{p_k \prod_{i=1, i\neq k}^{s}(p_i - p_k)} (k = 1, 2, \ldots, s)$ are positive integers. Then the complete r-partite graph $K_{p_1,p_2,\ldots,p_r} = K_{a_1\cdot p_1,\ldots,a_s\cdot p_s}$ is distance integral.

Theorem 3. *(See Corollary 2.9 of [16])For any positive integer q, the complete r-partite graph $K_{p_1q,p_2q,\ldots,p_rq} = K_{a_1\cdot p_1q,a_2\cdot p_2q,\ldots,a_s\cdot p_sq}$ is distance integral if and only if the complete r-partite graph $K_{p_1,p_2,\ldots,p_r=K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}}$ is distance integral.*

Remark 4. *Let $GCD(p_1, p_2, \ldots, p_s)$ denote the greatest common divisor of the numbers $p_1, p_2, \ldots, p_s$. We say that a vector $(p_1, p_2, \ldots, p_s)$ is primitive if $GCD(p_1, p_2, \ldots, p_s) = 1$. Theorem 3 shows that it is reasonable to study the complete r-partite graph $K_{p_1,p_2,\ldots,p_r} = K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$ only for primitive vectors $(p_1, p_2, \ldots, p_s)$.*

Theorem 5. *(See [16]) Let a complete r-partite graph $K_{p_1,p_2,\ldots,p_r} = K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$ be distance integral with eigenvalues μ_i. Let $\mu_i (\geq 0)$ and $p_i (> 0) (i = 1, 2, \ldots, s)$ be integers such that $-2 < p_1 - 2 < \mu_1 < p_2 - 2 < \mu_2 < \cdots < p_{s-1} - 2 < \mu_{s-1} < p_s - 2 < \mu_s < +\infty$ and*

$$a_k = \frac{\prod_{i=1}^{s}(\mu_i - p_k + 2)}{p_k \prod_{i=1, i\neq k}^{s}(p_i - p_k)}, k = 1, 2, \ldots, s \tag{3}$$

are positive integers, then for

$$b_k = \frac{\prod_{i=1}^{s-1}(\mu_i - p_k + 2)(\mu_s - p_k + 2 + rt)}{p_k \prod_{i=1,i\neq k}^{s}(p_i - p_k)}, k = 1, 2, \ldots, s, \tag{4}$$

$$r = LCM(r_1, r_2, \ldots, r_s), r_k = \frac{p_k \prod_{i=1,i\neq k}^{s}(p_i - p_k)}{d_k}, k = 1, 2, \ldots, s, \tag{5}$$

$$d_k = GCD(\prod_{i=1}^{s-1}(\mu_i - p_k + 2), p_k \prod_{i=1,i\neq k}^{s}(p_i - p_k)), k = 1, 2, \ldots, s. \tag{6}$$

the complete m-partite graph $K_{p_1,p_2,\ldots,p_m} = K_{b_1\cdot p_1, b_2\cdot p_2,\ldots,b_s\cdot p_s}$ *is distance integral for every non-negative integer* t *with eigenvalues* $\mu_1, \mu_2, \ldots, \mu_{s-1}, \mu_s' = \mu_s + rt$. *(Similar results for integral complete multipartite graphs were given in [5])*

3 Main Results

In this section, we construct some distance integral complete r-partite graphs with $s = 5, 6$. Moreover we also get infinitely many new classes of distance integral complete r-partite graphs $K_{p_1,p_2,\ldots,p_r} = K_{a_1\cdot p_1, a_2\cdot p_2,\ldots,a_s\cdot p_s}$ with $s = 5, 6$ by using Theorem 5.(Similar results for integral complete multipartite graphs were given in [14])

Theorem 6. *For* $s = 5$, *integers* $p_i(> 0)$, $a_i(> 0)$ *and* $\mu_i(i = 1, 2, 3, 4, 5)$ *are given in Table 1.* p_i, a_i *and* $\mu_i(i = 1, 2, 3, 4, 5)$ *are those of Theorem 2, where* $GCD(p_1, p_2, p_3, p_4, p_5) = 1$. *Then for any positive integer* q *the graph* $K_{a_1\cdot p_1 q, a_2\cdot p_2 q, a_3\cdot p_3 q, a_4\cdot p_4 q, a_5\cdot p_5 q}$ *on* $qn(= q\sum_{i=1}^{5} a_i p_i)$ *vertices is distance integral.*

Proof. From Theorem 2, it is not difficult to find that for $s = 5$ the complete r-partite graphs $K_{p_1,p_2,\ldots,p_r} = K_{a_1\cdot p_1, a_2\cdot p_2,\ldots,a_s\cdot p_s}$ is distance integral if and only if integers $p_i(> 0)$, $a_i(> 0)$ and $\mu_i(i = 1, 2, 3, 4, 5)$ satisfy the following equations.

$$a_1 = \frac{(\mu_1 - p_1 + 2)(\mu_2 - p_1 + 2)(\mu_3 - p_1 + 2)(\mu_4 - p_1 + 2)(\mu_5 - p_1 + 2)}{p_1(p_2 - p_1)(p_3 - p_1)(p_4 - p_1)(p_5 - p_1)},$$

$$a_2 = \frac{(\mu_1 - p_2 + 2)(\mu_2 - p_2 + 2)(\mu_3 - p_2 + 2)(\mu_4 - p_2 + 2)(\mu_5 - p_2 + 2)}{p_2(p_1 - p_2)(p_3 - p_2)(p_4 - p_2)(p_5 - p_2)},$$

$$a_3 = \frac{(\mu_1 - p_3 + 2)(\mu_2 - p_3 + 2)(\mu_3 - p_3 + 2)(\mu_4 - p_3 + 2)(\mu_5 - p_3 + 2)}{p_3(p_1 - p_3)(p_2 - p_3)(p_4 - p_3)(p_5 - p_3)},$$

$$a_4 = \frac{(\mu_1 - p_4 + 2)(\mu_2 - p_4 + 2)(\mu_3 - p_4 + 2)(\mu_4 - p_4 + 2)(\mu_5 - p_4 + 2)}{p_4(p_1 - p_4)(p_2 - p_4)(p_3 - p_4)(p_5 - p_4)},$$

$$a_5 = \frac{(\mu_1 - p_5 + 2)(\mu_2 - p_5 + 2)(\mu_3 - p_5 + 2)(\mu_4 - p_5 + 2)(\mu_5 - p_5 + 2)}{p_5(p_1 - p_5)(p_2 - p_5)(p_3 - p_5)(p_4 - p_5)},$$

From Theorem 3, we need only consider the case $GCD(p_1, p_2, \ldots, p_s) = 1$. Hence, by using a computer search, we have found 37 integral solutions listed in Table 1, where $1 \leq p_1 \leq 7$, $p_1 + 1 \leq p_2 \leq 15$, $p_2 + 1 \leq p_3 \leq 20$, $p_3 + 1 \leq p_4 \leq 24$, $p_4 + 1 \leq p_5 \leq 50$, $p_1 - 2 < \mu_1 < p_2 - 2$, $p_2 - 2 < \mu_2 < p_3 - 2$, $p_3 - 2 < \mu_3 < p_4 - 2$, $p_4 - 2 < \mu_4 < p_5 - 2$, $\mu_5 < 10000$, and $n = \sum_{i=1}^{5} a_i p_i$.

By Theorem 3 it follows that these graphs $K_{a_1\cdot p_1 q, a_2\cdot p_2 q, a_3\cdot p_3 q, a_4\cdot p_4 q, a_5\cdot p_5 q}$ are distance integral for any positive integer q. □

According to Theorem 3 and Theorem 5, we can obtain the following theorem.

Table 1: Distance integral complete r-partite graphs $K_{a_1 \cdot p_1, a_2 \cdot p_2, a_3 \cdot p_3, a_4 \cdot p_4, a_5 \cdot p_5}$.

p_1	p_2	p_3	p_4	p_5	a_1	a_2	a_3	a_4	a_5	μ_1	μ_2	μ_3	μ_4	μ_5
1	3	5	12	20	1302	254	185	70	132	0	2	7	13	6478
1	3	8	18	26	1914	520	197	36	46	0	4	14	21	6901
1	3	9	23	35	1596	444	91	112	22	0	5	13	31	7105
1	3	12	23	37	580	155	77	12	8	0	5	19	32	2551
1	4	8	18	31	3282	332	101	22	112	1	5	14	20	9298
1	4	13	18	33	2355	230	82	47	85	1	7	14	23	8006
1	5	8	22	36	754	20	47	24	24	2	4	13	27	2638
1	5	9	19	35	1099	127	82	32	64	1	5	13	23	5337
1	5	9	22	36	1688	84	30	57	16	2	6	13	31	4219
1	5	11	19	37	1933	299	18	116	51	1	8	11	29	7731
1	6	8	10	19	3112	157	106	136	100	2	5	7	14	8168
1	6	8	21	28	1915	232	184	21	155	1	5	14	20	9574
1	6	9	16	24	245	19	9	2	4	2	6	13	19	574
1	6	10	20	40	2436	56	74	45	56	3	6	14	28	6668
1	6	12	20	32	1638	83	98	68	56	2	6	14	25	6478
1	6	15	23	45	4256	143	82	36	32	3	10	19	37	8623
1	7	9	17	20	1071	33	31	18	42	3	6	11	16	2735
1	7	11	17	26	344	113	39	56	68	0	7	11	19	4299
1	7	15	17	26	2920	129	53	48	105	3	9	14	19	8175
1	8	10	18	23	1064	107	56	99	140	1	7	10	18	7496
1	8	13	19	33	3225	180	74	92	68	3	9	14	26	9631
1	8	13	22	31	1228	102	22	51	96	2	9	13	23	6446
1	8	13	22	31	1535	144	44	102	30	2	9	14	27	6446
1	8	13	22	31	1842	198	88	51	24	2	9	17	27	6446
1	8	16	21	39	3873	60	63	22	16	5	11	18	34	6454
1	15	20	24	36	986	29	6	22	27	6	16	19	28	3058
2	4	8	21	35	370	150	30	18	64	1	5	12	22	4218
2	5	9	20	34	387	30	79	56	21	2	4	12	28	3483
2	5	10	14	22	96	7	19	15	6	2	4	10	18	768
2	5	11	15	23	908	590	134	51	80	1	7	12	18	8853
2	6	10	15	35	1225	46	120	164	46	3	5	10	28	8008
2	6	10	15	35	2058	184	80	41	39	3	7	12	28	8008
2	6	14	23	32	413	20	28	70	18	3	6	14	28	3540
2	6	14	23	32	590	50	7	22	45	3	10	14	24	3540
2	7	14	24	40	564	60	147	131	54	2	6	16	33	8930
2	8	13	20	32	1036	183	24	58	72	3	10	14	24	7326
2	9	14	20	50	552	44	118	50	16	3	8	16	42	4968
2	10	14	19	22	258	81	52	7	33	2	10	16	18	2924
2	10	16	21	26	198	28	10	3	6	4	12	18	22	1064
2	10	16	23	35	654	258	63	57	62	2	12	18	28	8393
2	10	16	23	35	1962	86	42	45	54	6	12	18	28	8393
2	10	20	23	34	2575	133	47	30	52	6	15	20	28	9888
3	6	9	16	28	387	254	70	92	128	2	6	10	19	8386
3	6	9	16	28	516	381	140	23	104	2	6	13	19	8386
3	6	15	20	33	532	82	86	33	112	3	7	16	22	7753
3	6	15	20	33	798	164	43	24	98	3	10	16	22	7753
3	7	12	19	35	450	148	20	51	45	3	9	13	26	5185
3	12	16	23	31	411	51	13	3	46	6	13	19	22	3563
3	12	17	19	30	410	215	60	74	50	3	13	16	25	7750
4	7	12	19	40	78	51	40	21	31	3	7	14	26	2810
4	8	15	20	25	230	170	189	101	38	3	8	16	22	8098
4	9	14	20	48	179	70	12	81	9	4	10	13	42	3582
4	9	16	22	39	129	136	22	18	134	3	11	16	22	7742
5	8	13	22	31	379	85	55	30	80	5	9	16	23	6446
5	10	14	17	42	385	156	66	163	27	5	10	13	36	8328
5	11	19	22	35	344	249	24	75	21	5	15	18	31	7313

Theorem 7. *For $s = 5$, let $p_i(> 0)$, $a_i(> 0)$ and $\mu_i(> 0)$ $(i = 1, 2, 3, 4, 5)$ be those of Theorem 2. Then for any positive integer q, the complete multipartite graphs $K_{a_1\cdot p_1q, a_2\cdot p_2q, \ldots, a_5\cdot p_5q}$ with $qn(= q\sum_{i=1}^{5} a_ip_i)$ vertices are distance integral if $p_1 = 1$, $p_2 = 4$, $p_3 = 8$, $p_4 = 18$, $p_5 = 31$, $\mu_1 = 1$, $\mu_2 = 5$, $\mu_3 = 14$, $\mu_4 = 20$, $\mu_5 = 9298 + 13236132t$, $a_1 = 4671576t + 3282$, $a_2 = 472719t + 332$, $a_3 = 143871t + 101$, $a_4 = 31372t + 22$ $a_5 = 159936t + 112$, and $n = \sum_{i=1}^{5} a_ip_i = 324632t + 2622$, where t is a nonnegative integer.*

Proof. The proof directly follows from Theorem 3 and Theorem 5. □

Remark 8. *Analogously to Theorem 7 it is possible to find conditions for parameters p_i, $a_i(i = 1, 2, 3, 4, 5)$ which depend on t for each graph in Table 1. By this procedure we get new classes of distance integral graphs.*

Theorem 9. *For $s = 6$, let $p_1 = 2$, $p_2 = 5$, $p_3 = 11$, $p_4 = 18$, $p_5 = 21$ and $p_6 = 23$, $\mu_1 = 1$, $\mu_2 = 7$, $\mu_3 = 12$, $\mu_4 = 18$, $\mu_5 = 20$ and $\mu_6 = 53979$, $a_1 = 4735$, $a_2 = 2941$, $a_3 = 514$, $a_4 = 593$, $a_5 = 213$, $a_6 = 391$. Then for any positive integer q the graph $K_{a_1\cdot p_1q, a_2\cdot p_2q,\ a_3\cdot p_3q, a_4\cdot p_4q, a_5\cdot p_5q, a_6\cdot p_6q}$ on $qn(= q\sum_{i=1}^{6} a_ip_i)$ vertices is distance integral.*

Proof. From Theorem 2, it is not difficult to find that for $s = 6$ the complete r-partite graphs $K_{p_1,p_2,\ldots,p_r} = K_{a_1\cdot p_1, a_2\cdot p_2, \ldots, a_s\cdot p_s}$ is distance integral if and only if integers $p_i(> 0)$, $a_i(> 0)$ and $\mu_i(i = 1, 2, 3, 4, 5, 6)$ satisfy the following equations.

$$a_1 = \frac{(\mu_1 - p_1 + 2)(\mu_2 - p_1 + 2)(\mu_3 - p_1 + 2)(\mu_4 - p_1 + 2)(\mu_5 - p_1 + 2)(\mu_6 - p_1 + 2)}{p_1(p_2 - p_1)(p_3 - p_1)(p_4 - p_1)(p_5 - p_1)(p_6 - p_1)},$$

$$a_2 = \frac{(\mu_1 - p_2 + 2)(\mu_2 - p_2 + 2)(\mu_3 - p_2 + 2)(\mu_4 - p_2 + 2)(\mu_5 - p_2 + 2)(\mu_6 - p_2 + 2)}{p_2(p_1 - p_2)(p_3 - p_2)(p_4 - p_2)(p_5 - p_2)(p_6 - p_2)},$$

$$a_3 = \frac{(\mu_1 - p_3 + 2)(\mu_2 - p_3 + 2)(\mu_3 - p_3 + 2)(\mu_4 - p_3 + 2)(\mu_5 - p_3 + 2)(\mu_6 - p_3 + 2)}{p_3(p_1 - p_3)(p_2 - p_3)(p_4 - p_3)(p_5 - p_3)(p_6 - p_3)},$$

$$a_4 = \frac{(\mu_1 - p_4 + 2)(\mu_2 - p_4 + 2)(\mu_3 - p_4 + 2)(\mu_4 - p_4 + 2)(\mu_5 - p_4 + 2)(\mu_6 - p_4 + 2)}{p_4(p_1 - p_4)(p_2 - p_4)(p_3 - p_4)(p_5 - p_4)(p_6 - p_4)},$$

$$a_5 = \frac{(\mu_1 - p_5 + 2)(\mu_2 - p_5 + 2)(\mu_3 - p_5 + 2)(\mu_4 - p_5 + 2)(\mu_5 - p_5 + 2)(\mu_6 - p_5 + 2)}{p_5(p_1 - p_5)(p_2 - p_5)(p_3 - p_5)(p_4 - p_5)(p_6 - p_5)},$$

$$a_6 = \frac{(\mu_1 - p_6 + 2)(\mu_2 - p_6 + 2)(\mu_3 - p_6 + 2)(\mu_4 - p_6 + 2)(\mu_5 - p_6 + 2)(\mu_6 - p_6 + 2)}{p_6(p_1 - p_6)(p_2 - p_6)(p_3 - p_6)(p_4 - p_6)(p_5 - p_6)},$$

From Theorem 3, we need only consider the case $GCD(p_1, p_2, \ldots, p_s) = 1$. Hence, by using a computer search, we found this result, where $1 \le p_1 \le 5$, $p_1+1 \le p_2 \le p_1+10$, $p_2+1 \le p_3 \le p_2+10$, $p_3 + 1 \le p_4 \le p_3 + 10$, $p_4 + 1 \le p_5 \le p_4 + 10$, $p_5 + 1 \le p_6 \le p_5 + 20$, $p_1 - 2 < \mu_1 < p_2 - 2$, $p_2 - 2 < \mu_2 < p_3 - 2$, $p_3 - 2 < \mu_3 < p_4 - 2$, $p_4 - 2 < \mu_4 < p_5 - 2$, $p_5 - 2 < \mu_5 < p_6 - 2$, $p_6 - 2 < \mu_6 < 200000$ and $n = \sum_{i=1}^{6} a_ip_i$.

By Theorem 3 it follows that these graphs $K_{a_1\cdot p_1q, a_2\cdot p_2q, a_3\cdot p_3q, a_4\cdot p_4q, a_5\cdot p_5q, a_6\cdot p_6q}$ are distance integral for any positive integer q. □

According to Theorem 3 and Theorem 5, we can obtain the following theorem.

Theorem 10. *For $s = 6$, let $p_i(> 0)$, $a_i(> 0)$ and $\mu_i(> 0)$ $(i = 1, 2, 3, 4, 5, 6)$ be those of Theorem 2. Then for any positive integer q, the complete multipartite graphs $K_{a_1\cdot p_1q, a_2\cdot p_2q, a_3\cdot p_3q, \ldots, a_5\cdot p_5q, a_6\cdot p_6q}$ with $qn(= q\sum_{i=1}^{6} a_ip_i)$ vertices are distance integral if $p_1 = 2$, $p_2 = 5$, $p_3 = 11$, $p_4 = 18$, $p_5 = 21$, $p_6 = 23$, $\mu_1 = 1$, $\mu_2 = 7$, $\mu_3 = 12$, $\mu_4 = 18$, $\mu_5 = 20$, $\mu_6 = 9598038t + 53979$, $a_1 = 418600t+4735$, $a_2 = 260015t+2941$, $a_3 = 45448t+514$, $a_4 = 52440t+593$, $a_5 = 18837t+213$, $a_6 = 34580t + 391$, $n = \sum_{i=1}^{6} a_ip_i = 829920t + 9387$, where t is a nonnegative integer.*

Proof. The proof directly follows from Theorem 3 and Theorem 5. □

References

[1] AOUCHICHE M, HANSEN P. Two Laplacians for the distance matrix of a graph[J]. Linear Algebra Appl, 2013, 439: 21–33.

[2] BOSE S S, NATH M, PAUL S. Distance spectral radius of graphs with r pendent vertices[J]. Linear Algebra Appl, 2011, 435: 2828–2836.

[3] BUCKLEY F, HARARY F. Distance in Graphs[M]. Addison-Wesley, Redwood City, CA, 1990.

[4] DAS K C. On the largest eigenvalue of the distance matrix of a bipartite graph[J]. MATCH Commun Math Comput Chem, 2009, 62: 667–672.

[5] HÍC P, POKORNÝ M. New Classes of Integral Complete n-partite Graphs[J]. Advances and Applications in Discrete Mathematics, 2011, 7(2): 83–94.

[6] INDULAL G, GUTMAN I. On the distance spectra of some graphs[J]. Math Commun, 2008, 13: 123–131.

[7] LIN H Q, HONG Y, WANG J F, SHU J L. On the distance spectrum of graphs[J]. Linear Algebra Appl, 2013, 439: 1662–1669.

[8] LU S F, ZHAO H X, Signless Laplacian characteristic polynomials of complete multipartite graphs[J]. Chinese Quarterly Journal of Mathematics, 2012, 27(1): 36–40.

[9] MERRIS R. The distance spectrum of a tree[J]. J. Graph Theory, 1990, 14: 365–369.

[10] STEVANOVIĆ D, MILOSEVIĆ M, HÍC P, POKORNÝ M. Proof of a conjecture on distance energy of complete multipartite graphs[J]. MATCH Commun Math Comput Chem, 2013, 70: 157-162.

[11] STEVANOVIĆ D, ILIĆ A. Distance spectral radius of trees with fixed maximum degree[J]. Electron J Linear Algebra, 2010, 20: 168–179.

[12] STEVANOVIĆ D, INDULAL G. The distance spectrum and energy of the compositions of regular graphs[J]. Appl Math Lett, 2009, 22: 1136–1140.

[13] TAN S W, On the Laplacian spectral radius of trees[J], Chinese Quarterly Journal of Mathematics, 2010, 25(4): 615–625.

[14] WANG L G, WANG Q. Integral complete multipartite graphs $K_{a_1\cdot p_1,a_2\cdot p_2,\ldots,a_s\cdot p_s}$ with s=5,6[J]. Discrete Math, 2010, 310: 812–818.

[15] WANG Y N, ZHOU B. On distance spectral radius of graphs[J]. Linear Algebra Appl, 2013, 438: 3490–3503.

[16] YANG R S, WANG L G. Distance integral complete r-partite graphs[J]. Filomat, accepted, 2014.

[17] ZHOU B. On the largest eigenvalue of the distance matrix of a tree[J]. MATCH Commun Math Comput Chem, 2007, 58: 657–662.

[18] ZHOU B, ILIĆ A. On distance spectral radius and distance energy of graphs[J]. MATCH Commun Math Comput Chem, 2010, 64: 261–280.

[19] ZHOU H Q. The rank of integral circulant graphs[J]. Chinese Quarterly Journal of Mathematics, 2014, 29(1):116–124.